

Aberystwyth University

Enterobactin Deficiency in a Coliform Mastitis Isolate Decreases Its Fitness in a Murine Model

Vander Elst, Niels; Breyne, Koen; Steenbrugge, Jonas; Gibson, Amanda Jane; Smith, David George Emslie; Germon, Pierre; Werling, Dirk; Meyer, Evelyne

Published in:
Frontiers in Veterinary Science

DOI:
[10.3389/fvets.2020.576583](https://doi.org/10.3389/fvets.2020.576583)

Publication date:
2020

Citation for published version (APA):
Vander Elst, N., Breyne, K., Steenbrugge, J., Gibson, A. J., Smith, D. G. E., Germon, P., Werling, D., & Meyer, E. (2020). Enterobactin Deficiency in a Coliform Mastitis Isolate Decreases Its Fitness in a Murine Model: A Preliminary Host–Pathogen Interaction Study. *Frontiers in Veterinary Science*, 7, [576583].
<https://doi.org/10.3389/fvets.2020.576583>

Document License CC BY

General rights

Copyright and moral rights for the publications made accessible in the Aberystwyth Research Portal (the Institutional Repository) are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Aberystwyth Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Aberystwyth Research Portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

tel: +44 1970 62 2400
email: is@aber.ac.uk

Gene	Forward primer sequence 5'-3'	Reverse primer sequence 5'-3'
LCN2	TCTGTTGGAGGATGGAACTTCG	TAACAGGATGGAGGTGACGTTG
18s	CGGGGAGGTAGTGACGAAA	CCGCTCCCAAGATCCAATA
PPIA	TCCGGGATTTATGTGCCAGGG	GCTTGCCATCCAACCACTCAG

Supplemental Table 1: Primers used for RT-qPCR after reverse transcription of mRNA, derived 24 h.p.i. from bovine udder quarters experimentally infected with 500 CFU *E. coli* 1303.